






Hypertrophic Cardiomyopathy Panel

	Test Code	D4103
	Test Summary	This test analyzes 39 genes that have been associated with hypertrophic cardiomyopathy and disorders associated with hypertrophic cardiomyopathy.
	Turn-Around-Time (TAT)*	2 - 3 weeks
	Acceptable Sample Types	Whole Blood (EDTA) (Preferred sample type) DNA, Isolated Dried Blood Spots Saliva
	Acceptable Billing Types	Self (patient) Payment Institutional Billing Commercial Insurance

Indications for Testing

- Individuals with a clinical diagnosis of hypertrophic cardiomyopathy.
- Individuals with a known mutation in a gene that is associated with hypertrophic cardiomyopathy.

Test Description

This panel analyzes 39 genes that have been associated with hypertrophic cardiomyopathy and disorders associated with hypertrophic cardiomyopathy. Both sequencing and deletion/duplication (CNV) analysis will be performed on the coding regions of all genes included (unless otherwise marked). All analysis is performed utilizing Next Generation Sequencing (NGS) technology. CNV analysis detects most deletions and duplications of three exons or greater in size. All variants are classified according to ACMG guidelines.

Condition Description

Hypertrophic cardiomyopathy is a group of disorders characterized by the thickening of the myocardium. Symptoms vary even in the same family with variable expressivity; some individuals have no symptoms. Symptoms include but are not limited to heart murmurs, shortness of breath, palpitations, and chest pain. Familial hypertrophic cardiomyopathy is estimated to affect 1 in 500 worldwide.

Genes

A2ML1, ACADVL, ACTC1, ACTN2, AGL, ANKRD1, BAG3, BRAF, CACNA1C, CALR3, CAV3, CBL, CPT2, CSRP3, DES, ELAC2, FHL1, FLNC, GAA, GATA4, GLA, HRAS, JPH2, KRAS, LAMP2, LDB3, MAP2K1, MAP2K2, MTO1, MYBPC3, MYH6, MYH7, MYL2, MYL3, MYLK2, MYOM1, MYOZ2, MYPN, NEXN, NF1, NRAS, PDLIM3, PLN, PRKAG2, PTPN11, RAF1, RASA1, RIT1, RRAS, SHOC2, SOS1, SOS2, SPRED1, TCAP, TNNC1, TNNI3, TNNT2, TPM1, TTR, VCL, ACADVL, ACTC1, ACTN2, AGL, BRAF, CAV3, CBL, CSRP3, FHL1, FLNC, GAA, GLA, JPH2, LAMP2, MAP2K1, MYBPC3, MYH6, MYH7, MYL2, MYL3, MYOZ2, MYPN, NEXN, PLN, PRKAG2, RAF1, RIT1, SOS2, TCAP, TNNC1, TNNI3, TNNT2, TPM1, TTR, VCL, PTPN11, ALKP3, NRAS, SHOC2, SOS1

Test Methods and Limitations

Sequencing is performed on genomic DNA using an Agilent targeted sequence capture method to enrich for the genes on this panel. Direct sequencing of the amplified captured regions was performed using 2X100bp reads on Illumina next-generation sequencing (NGS) systems. A base is considered to have sufficient coverage at 20X, and an exon is considered fully covered if all coding bases plus three nucleotides of flanking sequence on either side are covered at 20X or more. Low coverage regions, if any, are limited to ~1% or less of the nucleotides in the test unless a pathogenic variant explaining the phenotype is discovered. A list of these regions is available upon request. Alignment to the human reference genome (hg19) is performed, and annotated variants are identified in the targeted region. Variants are called at a minimum coverage of 8X and an alternate allele

frequency of 20% or higher. Single nucleotide variants (SNVs) meeting internal quality assessment guidelines are confirmed by Sanger sequence analysis for records after results are reported. Indels and SNVs are confirmed by Sanger sequence analysis before reporting at the director's discretion. This assay cannot detect variants in regions of the exome that are not covered, such as deep intronic, promoter, and enhancer regions, areas containing large numbers of tandem repeats, and variants in mitochondrial DNA. Copy number variation (CNV) analysis detects deletions and duplications; in some instances, due to the size of the exons, sequencing complexity, or other factors, not all CNVs may be analyzed and/or may be difficult to detect. This assay is not designed to detect mosaicism; possible cases of mosaicism may be investigated at the discretion of the laboratory director. Primary data analysis is performed using Illumina DRAGEN Bio-IT Platform v.3.4.12. Secondary and tertiary data analysis is performed using PerkinElmer's internal ODIN v.1.01 software for SNVs and Biodiscovery's NxClinical v.6.1 or Illumina DRAGEN Bio-IT Platform v.3.4.12 for CNV and the absence of heterozygosity (AOH). Genes and/or exons located in pseudogene regions are not covered in this assay.

Detailed Sample Requirements

Whole Blood (EDTA) (Preferred sample type)

Collection Container(s):

EDTA (purple top)

Collection:

Infants (< 2-years): 2 to 3 mL; Children (>2-years): 3 to 5 mL; Older children and adults: Minimum 5mL. The blood tube should be inverted several times immediately after blood collection to prevent coagulation.

Sample Condition: Store at ambient temperature. Do not refrigerate or freeze.

Shipping: Ship overnight at ambient temperature ensuring receipt within 5-days of collection.

SPECIAL INSTRUCTIONS: Clotted or hemolyzed samples are not accepted.

DNA, Isolated

Collection:

Required DNA Quantity by Test Type*:

- **Next Generation Sequencing (NGS):** Send >1000 ng total gDNA @ >15 ng/?L. Please ship samples in 10mM Tris. Do not use EDTA.
- **Sanger Sequencing:** Send >500 ng total gDNA @ >15 ng/?L (varies by the size of the gene and the variants requested).
- **Non-Sanger Sequencing Tests:** Send >500 ng total gDNA @ >15 ng/?L.

Sample Condition: * Required DNA Quality: High molecular weight DNA (>12kb). A260/A280 reading should be ? 1.8. A260/230 a ratio range of 1.8 to 2.2. Contact the laboratory for specific amounts if total ng cannot be met.

Shipping: Ship overnight at ambient temperature.

SPECIAL INSTRUCTIONS:

- **Research Laboratories:** DNA extracted in research laboratories is not acceptable. Only under exceptional circumstances (e.g., proband not available) will DNA extracted in a research laboratory be accepted for clinical testing. Additional testing (e.g., of other family members) may be required to confirm results.
- **Laboratories outside the United States:** Non-US laboratories are not subject to CLIA regulations and will be reviewed on a case-by-case basis. Please call to speak with a laboratory genetic counselor prior to submitting a DNA sample from any non-CLIA certified laboratory.
- **Special Notes:** If extracted DNA is submitted, information regarding the method used for extraction should be sent along with the sample.



Dried Blood Spots

Saliva

Collection Container(s):

Oragene™ Saliva Collection Kit or ORAcollect-Dx kit

Collection:

Collect saliva on an Oragene™ Saliva Collection Kit ORAcollect-Dx kit according to the manufacturer's instructions.

Sample Condition: Store at ambient temperature. Do not refrigerate or freeze.

Shipping: Ship overnight at ambient temperature.

SPECIAL INSTRUCTIONS: Please contact PerkinElmer to request the saliva collection kit for patients that cannot provide a blood sample as whole blood is the preferred sample.